

Patient Name

DOB

Cell Phone

Address

INFLIXIMAB (Remicade & biosimilars)

Status	<input type="checkbox"/> New Therapy <input type="checkbox"/> Order Renewal <input type="checkbox"/> Dosage or Frequency Change			
Diagnosis	<input type="checkbox"/> ICD 10: K51.__ Ulcerative Colitis (UC)		<input type="checkbox"/> ICD 10: L40.9 Plaque Psoriasis	
	<input type="checkbox"/> ICD 10: K50.__ Crohn's Disease (CD)		<input type="checkbox"/> ICD 10: M45.__ Ankylosing Spondylitis (AS)	
	<input type="checkbox"/> ICD 10: M06.__ Rheumatoid Arthritis (RA)		<input type="checkbox"/> ICD 10: L40.5 Psoriatic Arthritis (PsA)	
	<input type="checkbox"/> ICD 10 Code: _____ Other: _____			
Pertinent Medical Hx	Patient's weight (most recent): _____ lbs / kg (circle one)			
	TB Status & Date: _____		Hepatitis B Status & Date: _____	
Labs	Labs to be drawn by: <input type="checkbox"/> Infusion Clinic <input type="checkbox"/> Referring Physician			
	If Infusion Clinic: _____			
Premeds	<input type="checkbox"/> No premeds necessary			
	<input type="checkbox"/> acetaminophen (Tylenol) <input type="checkbox"/> 500mg / <input type="checkbox"/> 650mg / <input type="checkbox"/> 1000mg PO			
	<input type="checkbox"/> cetirizine (Zyrtec) 10mg PO - OR - <input type="checkbox"/> diphenhydramine (Benadryl) 25mg IV			
	<input type="checkbox"/> methylprednisolone (Solu-Medrol) <input type="checkbox"/> 40mg / <input type="checkbox"/> 125mg IV			
	<input type="checkbox"/> other: _____			
IV Fluids	<input type="checkbox"/> NS TKO <input type="checkbox"/> Other: _____			
Medication Order	Drug: <input type="checkbox"/> Biosimilar or originator product, based on payer preference or Infusion Clinic preference (<i>preferred</i>) - OR - <input type="checkbox"/> Specific product: _____			
	Dose: <input type="checkbox"/> 3mg/kg <input type="checkbox"/> 5mg/kg <input type="checkbox"/> 7.5mg/kg <input type="checkbox"/> 10mg/kg <input type="checkbox"/> Other: _____			
	Dose will be rounded up to nearest 100mg - OR - <input type="checkbox"/> Give exact dose			
	Frequency: <input type="checkbox"/> Induction: Week 0, 2, 6, then every 8 weeks thereafter			
	<input type="checkbox"/> Maintenance: every 8 weeks. <input type="checkbox"/> other: _____			
	Refills: <input type="checkbox"/> x 1 year <input type="checkbox"/> x _____ doses <input type="checkbox"/> No refills; give this dose only.			
Monitoring	Monitor for signs/symptoms of hypersensitivity during infusion and 15 mins post-infusion. For any signs of infusion reaction: STOP infusion. Contact on-site provider for instruction.			
Additional Orders				
Physician Information	Physician Name	_____	NPI	_____
	Office Contact	_____	Phone	_____
	Provider Signature:	_____	Date	_____



Patient Name
DOB
Cell Phone
Address

INFLIXIMAB (Remicade & biosimilars)

REQUIRED DOCUMENTATION	
○ Patient Demographics & Insurance Information:	- Copy of patient's insurance card – front and back
○ Clinical / Progress Notes, supporting primary diagnosis:	- 2 most recent office notes - Medication history
○ Most Recent Labs:	- CMP and CBC - TB screening (PPD, QFT Gold or TSpot) - Baseline Liver Enzymes

FAX to (480) 400-6121
Intake Specialist (480) 927-3800
referral@infuseablecare.com